

COGNITIVE IMPAIRMENT PREDICTS THE OCCURRENCE OF HEPATIC ENCEPHALOPATHY AFTER TRANSJUGULAR INTRAHEPATIC PORTOSYSTEMIC SHUNT

Silvia Nardelli, Stefania Gioia, Chiara Pasquale, Iliaria Pentassuglio, Alessio Farcomeni^o, Manuela Merli, Filippo Maria Salvatori*, Leandra Nikolli, Sabrina Torrisi, Francesca Greco and Valeria Nicoletti and Oliviero Riggio.

Dept. of Clinical Medicine, Centre for the Diagnosis and Treatment of Portal Hypertension, “Sapienza” University of Rome, Rome, Italy

^oDept. of Public Health and Infectious Diseases, “Sapienza” University of Rome, Rome, Italy

*Dept. Of Radiology, “Sapienza” University of Rome, Rome, Italy

Short title: Prediction of post-TIPS hepatic encephalopathy

Key words: TIPS, hepatic encephalopathy, psychometric evaluation.

Abbreviations: PHES: Psychometric Hepatic Encephalopathy Score; DST: Digit-Symbol Test; TMT-A: Trail-Making Test A; TMT-B: Trail-Making Test B; SDT: Serial-Dotting Test; LTT: Line-Tracing Test; MHE: minimal hepatic encephalopathy; HE: hepatic encephalopathy; TIPS: transjugular intrahepatic porto-systemic shunt; sHR: subdistribution hazard ratio.

Disclosures: the authors declare that no conflict of interest exists concerning this paper

ABSTRACT

Introduction/Aim: Hepatic Encephalopathy (HE) is a major problem in patients treated with TIPS. The aim of the study was to establish whether pre-TIPS covert HE is an independent risk factor for the development of HE after TIPS.

Patients/Methods. 82 consecutive cirrhotic patients submitted to TIPS were included. All patients underwent the PHES to identify those affected by covert HE before a TIPS. The incidence of the first episode of HE was estimated, taking into account the nature of the competing risks in the data (death or liver transplantation). **Results.** Thirty-five (43%) patients developed overt HE. The difference of post TIPS HE was highly significant ($p=0.0003$) amongst patients with or without covert HE before a TIPS. 77 % of patients with post TIPS HE were classified as affected by covert HE before TIPS. Age: (sHR 1.05, CI 1.02-1.08, $p=0.002$); Child Pugh Score: (sHR 1.29, CI 1.06-1.56, $p=0.01$) and covert HE: (sHR 3.16 CI: 1.43-6.99 $p=0.004$) were associated with post TIPS HE. Taking into consideration only the results of PHES evaluation, the negative predicting value was 0.80 for all patients and 0.88 for the patients submitted to TIPS because of refractory ascites. Thus, a patient with refractory ascites, without covert HE before a TIPS, has almost 90% probability of being free of HE after TIPS. **Conclusions.** Psychometric evaluation before TIPS is able to identify most of the patients who will develop HE after a TIPS and can be used to select patients in order to have the lowest incidence of this important complication.

Abstract word count: 246

INTRODUCTION

Transjugular intrahepatic portosystemic shunt (TIPS) is currently used for the treatment of complications of portal hypertension, mainly variceal rebleeding and refractory ascites ^[1-9]. This procedure involves a major drawback: hepatic encephalopathy (HE). This complication has been reported in 30-55% ^[10-15] of cirrhotic patients within the first year and up to 10% of patients treated with a TIPS may experience a severe form of HE that is refractory to standard treatments and will need to be resolved by reducing the shunt diameter ^[13]. Unfortunately, no pharmacological treatment has yet proved able to reduce the incidence of post-TIPS HE. The only randomized controlled trial (RCT) carried out with this aim failed to show any beneficial effect of drugs commonly used in the treatment of HE ^[16]. More recently, a RCT was performed to assess the efficacy of polytetrafluoroethylene (PTFE) covered stents of different diameters (10 mm vs. 8 mm) on the incidence of post TIPS HE. Unfortunately, the trial was stopped because the stents with the smaller diameter were unable to control the complications of portal hypertension ^[17]. Thus, the selection of patients remains the only method to try to reduce the incidence of post TIPS HE and many studies have attempted to identify the predictors of this complication in order to select for TIPS only those patients with the lowest incidence. The factors identified as the most robust predictors of post TIPS HE were previous HE, age, a low porto-caval pressure gradient and a high Child Pugh score ^[13, 18-20]. Further factors were high creatinine levels ^[13] and low serum sodium concentration. ^[21] Despite the exclusion of patients with previous HE and advanced liver disease, the incidence of post TIPS HE still remains fairly high ^[22], thus other factors are involved.

In cirrhotic patients without TIPS, the presence of subclinical cognitive impairment, also known as covert HE ^[23], has been shown to be a strong predictor of the occurrence of overt HE ^[24]. Correspondingly, alterations in psychometric performance detected in a TIPS candidate before the procedure may help to identify the patients at risk of HE after the procedure. Actually, in our RCT ^[16], post TIPS HE developed more frequently in patients with an abnormal psychometric test (Trail Making Test A) before TIPS and, more recently, Berlioux P. et al. showed in 54 patients submitted

to a TIPS that the incidence of HE increased in those with abnormal critical flicker frequency before a TIPS [25]. However, in both studies the relationship between psychometric performance and post TIPS HE was present only at the univariate analysis and thus the role of cognitive impairment as a risk factor for HE after a TIPS remains uncertain.

The aim of the present study was to establish whether pre-TIPS covert HE is an independent risk factor for the development of HE and if the psychometric evaluation before a TIPS may be used for selecting patients in order to have the lowest rate of HE after a TIPS.

PATIENTS AND METHODS

From January 2011 to December 2014, all consecutive cirrhotic patients undergoing TIPS were considered eligible for the study. In our Centre, exclusion criteria for TIPS placement are: age >75 years; bilirubin levels >5 mg/dl; creatinine levels >3 mg/dl; a serious cardiac or pulmonary dysfunction; a Child-Pugh's score >11 (except for patients who were candidates for early TIPS); a model end-stage liver disease (MELD) score >18; the presence of portal thrombosis; a diagnosis of hepatic carcinoma; sepsis; spontaneous bacterial peritonitis. Present HE or previous spontaneous/recurrent HE is also a contraindication to TIPS; however, patients with only one episode of HE precipitated by variceal bleeding and ameliorated after the bleeding was controlled are not excluded. Other exclusion criteria were: alcohol/psychoactive drugs intake (positive alcoholaemia and/or benzodiazepines or opioid urine metabolites) at the moment of evaluation, unrelated neurological disease including dementia (mini mental state < 26), lack of compliance with psychometric evaluation because of language barriers or reduced visual acuity.

The purpose of the study, the enrolment and the details of the TIPS operation were clearly explained to all the patients before obtaining their written informed consent. The "Sapienza"

University of Rome Ethical Committee approved the collection of data of the patients for prognostic studies (Rif.1720/01.10.09).

All TIPS procedures were carried out by the same radiology team, using PTFE-covered stents of 10 mm diameter. The anesthesiological procedure ^[4, 26] and the technical details of TIPS with PTFE-covered stent-grafts implantation were previously described ^[27-28]. All the subjects were evaluated and followed by the same medical team by a prospective protocolled diagnostic work-up and a surveillance strategy.

The day before the procedure, a basal evaluation of HE, including an examination and grading of the patients' mental state, asterixis, and psychometric performance as well as the determination of venous blood ammonia were carried out. The evaluation of the degree of HE was based on the alteration of the patient's mental state using modifications of the West Haven Criteria ^[29]. The mental state was assessed in each patient by the same investigator using standardized tests and questions, as previously described ^[30]. All patients also underwent the psychometric hepatic encephalopathy score (PHES) battery of tests, including the digit-symbol-test (DST), trail-making-test A (TMT-A) and B (TMT-B), the serial-dotting-test (SDT) and the line-tracing-test (LTT). Each test was scored against age and education-adjusted norms for the Italian population. The PHES is the sum of integer scores of each test computed from the adjusted Z values, as follows: score = -3 for $Z \leq -3$, score -2 for $-3 < Z \leq -2$, score -1 for $-2 < Z \leq -1$, score 0 for $-1 < Z < 1$, score 1 for $Z \geq 1$. The PHES ≤ -4 was considered abnormal ^[30]. Blood samples from a peripheral vein were collected in iced tubes for the determination of ammonia, which was performed immediately after using the Ammonia Checker II (Menarini, Florence, Italy), as previously described ^[31].

None of the patients received any pharmacological treatment to prevent the occurrence of HE. After TIPS, the patients remained hospitalized for one week and then were followed up once a week in the outpatients department for the first month. The patients were then seen every 3 months and also contacted by phone every month for the first six months. Thereafter, the patients were seen every 6 months. Moreover, both the patients and their families were instructed about the importance

of an immediate contact with the medical staff should any alteration in their mental state occur between the scheduled visits. In particular, the family was instructed to refer the occurrence of lethargy, apathy, obvious personality changes, inappropriate behaviour or disorientation to time and space (corresponding to a grade-II alteration of the patients' mental state). In this case, the HE evaluation, including the psychometric performance, was repeated to confirm and stage the degree of HE. A grade II HE or higher was considered an episode of overt HE ^[23] and the patients were censored as HE+ patients. The occurrence of a recurrent HE (defined as at least three episodes of non precipitant-induced severe encephalopathy requiring hospitalization in the last three months despite continuous treatment with non-absorbable disaccharides) or a persistent HE (defined as the presence of a continuously detectable altered mental state with further episodic deterioration despite protein restriction and treatment with non-absorbable disaccharides) was also recorded and the patients were considered affected by refractory HE. Those patients with an overt episode of HE were then managed either as in- or out-patients, depending on the severity of the HE episode. Once developed, HE was treated with the oral administration of non-absorbable disaccharides or non-absorbable antibiotics. All potential HE precipitating events were treated and, when possible, avoided. The patients' outcomes considered for the statistical analysis were the first episode of HE, liver transplantation (LT) and death.

Statistical analysis

The data are reported as mean \pm SD. Comparisons between groups were performed by unpaired Student's t-test or chi-square. We estimated the cumulative incidence of the first episode of HE during the first six months of follow up taking into account the nature of the competing risks in the data (HE before LT, death or LT are competing events). As the study is estimating outcomes other than all- cause mortality a method based on multistate disease models was selected. The usual Cox regression model in this context might be severely biased ^[32] and the sub-distribution model of Fine and Gray was selected. The conditional sub-distribution hazard at multivariate analysis was

evaluated using the model of Fine and Gray ^[33]. We therefore report on the sub-distribution hazard ratios (sHR) rather than the usual HR, but the former have similar interpretations to the latter. The factors associated with the development of HE were initially evaluated by univariate models (using univariate Fine and Gray models) and then included in a multivariate analysis (according to multivariate Fine and Gray models). The final multivariate model was chosen in a forward fashion by minimizing the Bayesian Information Criterion.

In order to build a score for the prediction of HE post TIPS we built an event-history analysis model allowing for competing risks, therefore predicting the risk of the event. The maximum likelihood coefficients of the optimal model were used as weights for the new score. A time-dependent ROC curve ^[34] for censored data at 6 months of follow up was estimated using the NN estimation method, whilst significance tests and confidence intervals were assessed through the non-parametric bootstrap. Sensitivity, specificity, PPV and NPV were based on the estimated ROC at 6 months. Software R version 3.0.2 was used for all computations.

RESULTS

The demographic, clinical and biochemical characteristics of the patients are reported in Table 1.

During the whole follow up 35 of the 82 (43%) patients developed at least one episode of overt HE. In 3 of them, HE persisted and was refractory to standard treatment, needing the reduction of the stent calibre to ameliorate. This procedure was carried out 5, 6 and 8 months after TIPS and led to the resolution of HE symptoms in all patients. During the whole follow up, 13 patients died and 5 were transplanted.

The comparison between the 35 patients with overt HE and the 47 patients who did not develop HE after a TIPS is reported in Table 2. At the time of the TIPS placement, there were no

significant differences between the two groups in gender, aetiology and severity of liver disease (Child-Pugh and MELD score) and most biochemical parameters. However, age (55 vs 62 years) was significantly higher and serum sodium (138 vs 135 mEq/ml) significantly lower in patients with post TIPS HE. The porto-systemic gradient measured immediately after the shunt opening was 6.9 ± 3.9 in the HE patients group and 5.9 ± 3.2 in the patients who did not develop HE after a TIPS (NS). In the group of patients with HE after a TIPS, there was a higher prevalence of patients in whom TIPS was indicated because of refractory ascites, but the difference did not reach statistical significance ($p=0.08$). As far as the pre TIPS evaluation of the patients' cognitive function is concerned, according to our exclusion criteria, no patients had signs of HE at inclusion and only a few of them (11 patients, 13%) experienced one episode of precipitated HE before a TIPS.

77 % of the patients with post TIPS HE were classified as affected by covert HE before a TIPS according to the PHES evaluation. The corresponding figure in the group without HE was 32% and the difference was highly significant ($p=0.0005$). The cumulative incidence of HE after TIPS, taking into consideration LT and death as risks competing with HE development, is reported in Figure 1. The difference in the incidence of post TIPS HE was highly significant ($p=0.0003$) among patients with or without covert HE detected by PHES before a TIPS. This difference, however, was at the limit of statistical significance, if only the 37 patients operated because of variceal bleeding are considered (58% vs 28%; $p=0.07$), whereas it was highly significant in the 45 patients submitted to a TIPS because of refractory ascites (87% vs 36%; $p=0.0001$). Moreover, all the 3 patients with refractory HE belonged to the group of ascitic patients.

Age, severity of liver disease (Child- Pugh/ MELD), TIPS indication (varices/ascites) and covert HE were included in a Fine and Gray multivariate model analysis. Age: (sHR 1.05, CI 1.02-1.08, $p=0.0022$); C-P.SCORE: (sHR 1.29, CI 1.06-1.56, $p=0.0110$) and covert HE: (sHR 3.16 CI: 1.43-6.99 $p=0.0045$) were independently associated with post TIPS HE development. On the basis of these results a model was developed as follows: $\text{Age}/10 + \text{CP.SCORE} + 4.88$ if covert HE is present

with an AUC of 0.75. With a cut off of 17, sensitivity was 0.77 (0.72-0.82), specificity 0.75 (0.70-0.80), PPV 0.64 (0.59-0.69) and NPV 0.83 (0.79-0.87). If age and the C-P score are excluded and only the presence of covert HE before a TIPS is considered, the corresponding values are: sensitivity 0.74 (0.69-0.79), specificity 0.63 (0.58-0.68), PPV 0.55 (0.50-0.60) and NPV 0.80 (0.76-0.84). The NPV was 0.78 (0.71-0.85) in the group of patients submitted to TIPS to prevent variceal rebleeding and 0.88 (0.83-0.93) in patients operated on because of refractory ascites.

DISCUSSION

The identification of the risk factors for the development of HE after a TIPS is a relevant problem. In fact, post TIPS HE is very frequent and, although in a minority of patients, it may be persistent and refractory to medical treatment, thus affecting deeply the patients' quality of life. The problem is, in our opinion, particularly relevant in patients submitted to a TIPS because of refractory ascites. In fact, at variance with the TIPS performed for the prevention of variceal rebleeding, which can be life saving and without therapeutic alternatives, a patient with refractory ascites may be treated with a TIPS or repeated large volume paracentesis with a fairly similar efficacy at least in terms of survival. The lack of preventive measures, whether pharmacological or based on the optimization of the porto-systemic gradient reached after the procedure, makes the optimal selection of patients particularly crucial. Previous HE with the exception of that precipitated by variceal bleeding, especially if recidivant, aging and advanced liver failure are the most robust risk factors for post TIPS HE ^[35] and today most protocols and clinical studies consider the presence of these factors as a contraindication to TIPS. Nevertheless, the incidence of post TIPS HE continues to be regrettably high, suggesting that other factors may be important.

The working hypothesis of the present paper was that the subclinical cognitive impairment also known as covert HE, may be a predictive factor for overt HE development after a TIPS. This hypothesis is supported by the fact that covert HE is one of the strongest predictors of the occur-

rence of overt HE in the follow up in cirrhotic patients ^[24] and by the observation that TMT-A, one of the tests included in the PHES, the standard for the identification of patients affected by covert HE, as well as the critical flicker frequency, was significantly correlated with post TIPS HE at least at univariate analysis. Our results support the hypothesis, as the incidence of post TIPS HE was significantly different amongst patients with or without covert HE before a TIPS. Our observation is methodologically solid because it was obtained in a group of patients already selected for TIPS on the basis of the known risk factors for HE development (see inclusion criteria). Moreover, post TIPS HE was detected using the PHES, which is considered the standard method ^[23] and statistically analysed taking into consideration the risks competing with HE during the patients' follow up ^[36], such as death and liver transplantation. Finally, at variance with previous observations, covert HE before a TIPS was a predictor independently of a number of clinical and laboratory variables, age and the Child Pugh score at the multivariate analysis. The model derived from the results (including age, Child Pugh score and covert HE) showed a fairly good sensitivity and specificity in identifying patients with overt HE after a TIPS. On the basis of the above results we tried to calculate the possible applicability of the detection of covert HE before a TIPS as a criterion for the selection of patients. If only the results of the PHES evaluation (i.e. not considering age and Child Pugh score) are taken into consideration, the negative predicting value was 0.80 for all patients and 0.88 for the patients submitted to TIPS because of refractory ascites. This means that a patient with refractory ascites, without covert HE according to PHES before a TIPS, has almost 90% probability of being free of HE after a TIPS. In our opinion this observation may help to choose the use of TIPS in patients particularly susceptible to severe HE and with treatment alternatives.

A limitation of the present observation is inherent to the use of the PHES which is based on 5 paper and pencil psychometric tests and on Z-scores obtained in a reference population that are available only in Germany, Italy, Spain, India and Korea. Another limitation of our study is that we were not able to identify the risk factors for refractory HE, which is the most relevant problem in

these patients. This was due to the fact that, fortunately, only 3 patients developed this complication during the follow up, limiting the possibility to analyze these data statistically.

In conclusion, the psychometric evaluation before a TIPS is able to identify most of the patients who will develop HE after a TIPS and can be useful to select patients in order to have the lowest incidence of this important complication.

References

- 1) Cabrera J, Maynar M, Granados R, et al. Transjugular intrahepatic portosystemic shunt versus sclerotherapy in the elective treatment of variceal hemorrhage. *Gastroenterology* 1996; 110: 832-839.
- 2) Sanjal AJ, Freedman AM, Luketic VA, et al. Transjugular intrahepatic portosystemic shunts for patients with active variceal hemorrhage unresponsive to sclerotherapy. *Gastroenterology* 1996; 111: 138-146.
- 3) Rossle M, Deibert P, Haag K, et al. Randomised trial of transjugular-intrahepatic-portosystemic shunt versus endoscopy plus propranolol for prevention of variceal rebleeding. *Lancet* 1997; 349: 1043-1049.
- 4) Merli M, Salerno F, Riggio O, et al. Transjugular intrahepatic portosystemic shunt versus endoscopic sclerotherapy for the prevention of variceal bleeding in cirrhosis: a randomized multicenter trial. *Hepatology* 1998; 27: 40-45.
- 5) Lebrec D, Giuily N, Hadengue A, et al. Transjugular intrahepatic portosystemic shunts: comparison with paracentesis in patient with cirrhosis and refractory ascites: a randomized trial. *J Hepatol* 1996; 25: 135-144.
- 6) Rossle M, Ochs A, Gulberg V, et al. A comparison of paracentesis and transjugular intrahepatic portosystemic shunting in patients with ascites. *N Engl J Med* 2000; 342: 1701-1707.

- 7) Gines P, Uriz J, Calahorra B, et al. Transjugular intrahepatic portosystemic shunt versus paracentesis plus albumin for refractory ascites in cirrhosis. *Gastroenterology* 2002;123:1839-1847.
- 8) Sanyal AJ, Genning C, Reddy KR, et al. The North American study for the treatment of refractory ascites. *Gastroenterology* 2003; 124: 634-641.
- 9) Salerno F, Merli M, Riggio O, et al. Randomized controlled study of TIPS versus paracentesis plus albumin in cirrhosis with severe ascites. *Hepatology* 2004; 40: 629-635.
- 10) Sanjal AJ, Freedman A, Shiffman ML, et al. Portosystemic encephalopathy after transjugular intrahepatic portosystemic shunt: results of a prospective controlled study. *Hepatology* 1994; 20: 46-55.
- 11) Jalan R, Elton RA, Redhead DN, et al. Analysis of prognostic variables in the prediction of mortality, shunt failure, variceal rebleeding and encephalopathy following the Transjugular intrahepatic portosystemic shunt for variceal haemorrhage. *Journal of Hepatology* 1995; 23: 123-128.
- 12) Somberg KA, Riegler JL, LaBerge JM, et al. Hepatic encephalopathy after transjugular intrahepatic portosystemic shunt: incidence and risk factors. *Am J Gastroenterol* 1995; 90: 549-555.
- 13) Riggio O, Angeloni S, Salvatori FM, et al. Incidence, Natural History, and Risk Factors of Hepatic Encephalopathy After Transjugular Intrahepatic Portosystemic Shunt With Polytetrafluoroethylene-Covered Stent Grafts *Am J Gastroenterol* 2008;103:2738–2746
- 14) Zuckerman DA, Darcy MD, Bocchini TP, et al. Encephalopathy after transjugular intrahepatic portosystemic shunt: analysis and incidence of potential risk factors. *American Journal of Radiology*; 169: 1727-1731.


- 15) Nolte W, Wiltfang J, Schindler C, et al. Portosystemic hepatic encephalopathy after transjugular intrahepatic portosystemic shunt in patients with cirrhosis: clinical, laboratory, psychometric and electroencephalographic investigations. *Hepatology* 1998;28:1215-1225.
- 16) Riggio O, Masini A, Efrati C, et al. Pharmacological prophylaxis of hepatic encephalopathy after transjugular intrahepatic portosystemic shunt: a randomized controlled study. *J Hepatol* 2005; 42: 674-679.
- 17) Riggio O, Ridola L, Angeloni S et al. Clinical efficacy of transjugular intrahepatic portosystemic shunt created with covered stents with different diameters: results of a randomized controlled trial. *J Hepatol*. 2010 ;53:267-72.
- 18) Bai M, Qi X, Yang Z, et al. Predictors of hepatic encephalopathy after transjugular intrahepatic portosystemic shunt in cirrhotic patients: A systematic review *Journal of Gastroenterology and Hepatology* 2011; 26:943–951.
- 19) Rossle M, Deibert P, Haag K, et al. Randomised trial of transjugular-intrahepaticportosystemicshunt versus endoscopy plus propranolol for prevention of variceal rebleeding. *Lancet* 1997;349:1043–9.
- 20) Casado M, Bosch J, Garcia-Pagan JC et al. Clinical events after TIPS: correlation with hemodynamic findings. *Gastroenterology* 1998;114: 1296-1303.
- 21) Guevara M1, Baccaro ME, Ríos J, Martín-Llahí M, Uriz J, Ruiz del Arbol L, Planas R, Monescillo A, Guarner C, Crespo J, Bañares R, Arroyo V, Ginès P. Risk factors for hepatic encephalopathy in patients with cirrhosis and refractory ascites: relevance of serum sodium concentration. *Liver Int*. 2010 Sep; 30(8):1137-42.

- 22) Riggio O, Nardelli S, Pasquale C, Pentassuglio I, Gioia S, Onori E, Frieri C, Salvatori FM, Merli M. No effect of albumin infusion on the prevention of hepatic encephalopathy after transjugular intrahepatic portosystemic shunt. *Metab Brain Dis.* 2015 Aug 20. [Epub ahead of print]
- 23) Vilstrup H, Amodio P, Bajaj J, Cordoba J, Ferenci P, Mullen KD, Weissenborn K, Wong P. Hepatic encephalopathy in chronic liver disease: 2014 Practice Guideline by the American Association for the Study of Liver Diseases and the European Association for the Study of the Liver. *Hepatology.* 2014 Aug;60(2):715-35.
- 24) Riggio O, Amodio P, Farcomeni A, Merli M, Nardelli S, Pasquale C, Pentassuglio I, Gioia S, Onori E, Piazza N, De Rui M, Schiff S, Montagnese S. A Model for Predicting Development of Overt Hepatic Encephalopathy in Patients With Cirrhosis. *Clin Gastroenterol Hepatol.* 2015 Jul;13(7):1346-52.
- 25) Berlioux P, Robic MA, Poirson H, Métivier S, Otal P, Barret C, Lopez F, Péron JM, Vinel JP, Bureau C. Pre-transjugular intrahepatic portosystemic shunts (TIPS) prediction of post-TIPS overt hepatic encephalopathy: the critical flicker frequency is more accurate than psychometric tests. *Hepatology.* 2014 Feb;59(2):622-9.
- 26) Sampietro G, Rossi P, Di Marco P. Use of a laryngeal mask in transjugular intrahepatic portosystemic shunt procedures. *J Vasc Interv Radiol* 1998; 9: 169.
- 27) Angeloni S, Merli M, Salvatori F, et al. Polytetrafluorethylene-covered stent-graft for TIPS procedure: 1-year patency and clinical results. *Am J Gastroenterol* 2004; 99:280-285.
- 28) Rossi P, Salvatori F, Fanelli F, et al. Polytetrafluorethylene-covered nitinol stent-graft for transjugular intrahepatic portosystemic shunt creation: 3-year experience. *Radiology* 2004; 231: 820-830.

- 29) Conn HO, Leevy CM, Vlahcevic ZR, et al. Comparison of lactulose and neomycin in the treatment of chronic portal-systemic encephalopathy. A double blind controlled trial. *Gastroenterology* 1977; 72: 573-583.
- 30) Amodio P, Campagna F, Olinas S et al. Detection of minimal hepatic encephalopathy: normalization and optimization of the Psychometric Hepatic Encephalopathy Score. A neuro-psychological and quantified EEG study. *J Hepatol*. 2008; 49: 346-53.
- 31) Nicolao F, Efrati C, Masini A, Merli M, Attili AF, Riggio O. Role of determination of partial pressure of ammonia in cirrhotic patients with and without hepatic encephalopathy. *J Hepatol* 2003; 38: 441-446.
- 32) [Jepsen P](#), [Vilstrup H](#), [Andersen PK](#). The clinical course of cirrhosis: the importance of multistate models and competing risks analysis. [Hepatology](#). 2015 Jul;62(1):292-302.
- 33) Fine JP, Gray RJ. A proportional hazards model for the subdistribution of a competing risk. *J Am Stat Assoc* 1999;94:496–509.
- 34) Heagerty PJ, Lumley T, Pepe MS. (2000) Time-dependent ROC Curves for Censored Survival Data and a Diagnostic Marker. *Biometrics*, 56, 337 – 344.
- 35) Bai M, Qi X, Yang Z, Yin Z, Nie Y, Yuan S, Wu K, Han G, Fan D. Predictors of hepatic encephalopathy after transjugular intrahepatic portosystemic shunt in cirrhotic patients: a systematic review. *J Gastroenterol Hepatol*. 2011 Jun;26(6):943-51. doi: 10.1111/j.1440-1746.2011.06663.x. Review.
- 36) Jepsen P, Vilstrup H, Andersen PK. The clinical course of cirrhosis: The importance of multistate models and competing risks analysis. *Hepatology*. 2015 Jul;62(1):292-302.

Figure 1

"Cumulative incidence of HE post TIPS for patients with and without MHE, taking into account two additional competing risks: OLT and Death"



Patients
at risk

42	27	24	23	21	21	20
40	37	36	36	36	36	35

